


# Bottomonium Decays at CLEO

Jianchun Wang  
*Syracuse University*

*DPF 2006+JPS 2006, 10/31/06*

# The Detector and Upsilon Data Sets



10/31/06

Jianchun Wang


## CLEO III

Excellent Calorimeter Coverage and Resolution

Excellent Tracking Coverage and Resolution

Ring Imaging Cerenkov and  $dE/dx$  for PID

Muon Chambers



Great Accelerator Team **CESR**

Dedicated  $\Upsilon(1S)$ ,  $\Upsilon(2S)$ ,  $\Upsilon(3S)$  runs  
(Nov 2001 – Dec 2002).

# CLEO III Upsilon Topics

$$\Upsilon(4S) \rightarrow B\bar{B}$$

$$\Upsilon(nS) \rightarrow ggg, gg\gamma, q\bar{q}$$

$$\Upsilon(nS) \rightarrow \Upsilon(mS)X, \chi_b X, \dots$$



$$e^+e^- \rightarrow q\bar{q}$$

$$\Upsilon(nS) \rightarrow ee, \mu\mu, \tau\tau$$

- ◆ Anti-deuteron production in  $\Upsilon(nS)$  decays and the nearby continuum.
- ◆ Inclusive production  $\Upsilon(1S) \rightarrow \eta' X$ .
- ◆ Upsilon radiative decays:
  - Observation of exclusive modes:  $\gamma\pi\pi, \gamma KK$ .
  - Search for  $\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0, \gamma\eta\eta, \gamma\pi^0\eta$ .
  - Search for  $\Upsilon(1S) \rightarrow \gamma\eta, \gamma\eta'$ .
  - UL on multi-body modes ( $\geq 4$  charged tracks).
  - Comparison of inclusive hadron production in gluon-rich and quark environment.

# Anti-Deuteron in Upsilon Decays

- ◆ Anti-deuteron production has been observed in many different kinds of hadronization processes.
- ◆ ARGUS observed anti-deuteron in  $\Upsilon(1S)$  decays. OPAL set an upper limit in Z decays.
- ◆ The production can be explained by coalescence model and a model based on string calculation. The basic idea is that an anti-proton and an anti-neutron are produced nearby in phase space to form an anti-deuteron.
- ◆ CLEO has a factor  $\sim 40$  more  $\Upsilon(1S)$  than ARGUS to make a more precise measurement.
- ◆ The main PID tool is  $dE/dX$  with RICH info also used.
- ◆ CLEO measured production in momentum range (0.45 – 1.45) GeV/c and use a model dependent extrapolation for unmeasured region.



$$\chi_d \equiv \frac{(dE/dx)_{\text{measured}} - (dE/dx)_{\text{expected},d}}{\sigma_{dE/dx}}$$

10/31/06

Jianchun Wang

# Anti-Deuteron Production

- ◆ The production BR per direct  $\Upsilon(1S) \rightarrow ggg, g\bar{g}$  decays is

$$B^{dir}(\Upsilon(1S) \rightarrow \bar{d}X) = (3.36 \pm 0.23 \pm 0.25) \times 10^{-5}$$

The overall BR per  $\Upsilon(1S)$  is

$$B(\Upsilon(1S) \rightarrow \bar{d}X) = (2.86 \pm 0.19 \pm 0.21) \times 10^{-5}$$

- ◆ The production in  $\Upsilon(2S)$  is

$$B(\Upsilon(2S) \rightarrow \bar{d}X) = (3.37 \pm 0.50 \pm 0.25) \times 10^{-5}$$

Removing contribution of  $\Upsilon(2S) \rightarrow \Upsilon(1S)X$ , and  $\Upsilon(2S) \rightarrow ggg, g\bar{g}, q\bar{q}$ , CLEO set 90% CL limit


$$B(g_{b_{1,2,0}} \rightarrow \bar{d}X) < 1.09 \times 10^{-4}$$

- ◆ The production rate upper limit in  $\Upsilon(4S)$  is

$$B(\Upsilon(4S) \rightarrow \bar{d}X) < 1.31 \times 10^{-5}$$

- ◆ Continuum production cross section at  $\sqrt{s}=10.5$  GeV

$$B(e^+e^- \rightarrow \bar{d}X) < 0.031 \text{ pb}$$




Deuteron production is enhanced in  $ggg, g\bar{g}$  process.

# Inclusive $\Upsilon(1S) \rightarrow \eta' X$

- ◆ The unexpectedly large  $B(B \rightarrow \eta' X_S)$  with  $P_{\eta'} > 2$  GeV was observed by CLEO and confirmed by BaBar.

$$\begin{aligned} \text{CLEO} &: (6.2 \pm 1.6 \pm 1.3^{+0.0}_{-1.5}) \times 10^{-4} \\ \text{BaBar} &: (3.9 \pm 0.8 \pm 0.5 \pm 0.8) \times 10^{-4} \end{aligned}$$

- ◆ Within SM, a possible explanation is the large  $g^* \rightarrow g\eta'$  coupling in  $b \rightarrow sg$  penguin diagram (*Atwood & Soni*).




- ◆ Such enhancement should present in  $\Upsilon(1S)$  decays (*Kagan, Atwood & Soni*).
- ◆ CLEO II measured  $B(\Upsilon(1S) \rightarrow ggg \rightarrow \eta' X) = (1.9 \pm 1.1 \pm 0.2) \times 10^{-4}$  for  $E_{\eta'}/E_{\text{beam}} > 0.7$ , ruled out a class of form factors characterized by a weak  $q^2$  dependence. Higher precision is needed for more model tests.

# Inclusive $\eta'$ Production In $\Upsilon(1S)$ Data

$\eta' \rightarrow \eta\pi^+\pi^-, \eta \rightarrow \gamma\gamma$

	All Z	Z > 0.7
a) $n \equiv \frac{N(\Upsilon(1S) \rightarrow ggg \rightarrow \mathbf{h}' X)}{N(\Upsilon(1S) \rightarrow ggg)}$	$(3.2 \pm 0.2 \pm 0.2)\%$	$(3.7 \pm 0.5 \pm 0.3) \times 10^{-4}$
b) $n \equiv \frac{N(\Upsilon(1S) \rightarrow q\bar{q} \rightarrow \mathbf{h}' X)}{N(\Upsilon(1S) \rightarrow q\bar{q})}$	$(3.8 \pm 0.2 \pm 0.3)\%$	$(22.5 \pm 1.2 \pm 1.8) \times 10^{-4}$
c) $n \equiv \frac{N(\Upsilon(1S) \rightarrow \mathbf{h}' X)}{N(\Upsilon(1S))}$	$(3.0 \pm 0.2 \pm 0.2)\%$	$(5.1 \pm 0.4 \pm 0.4) \times 10^{-4}$




10/31/06

Jianchun Wang

7

# Inclusive $\eta'$ Production Compare with Theory




- ◆ Measured spectrum is not well described by existing models.
- ◆ The observed  $B(B \rightarrow \eta' X_S)$  is unlikely to be explained by an enhanced  $g^* g \eta'$  form factor. An explanation outside the realm of SM or an improved understanding of non-perturbative QCD effects may be needed.




# Upsilon Radiative Decays

- ◆ In Upsilon two-body radiative decays, The two gluons hadronize into a meson or form a glueball.



- ◆ These decays are theoretically simple (no hadronic FSI). They are useful in study of color-singlet two-gluon system.
- ◆ Many  $J/\psi \rightarrow \gamma X$  ( $X=\eta, \eta', \eta_c, f_2(1270), \dots$ ) decay modes have been observed. A possible glueball state  $f_J(2220)$  was reported by BES. Other candidates like  $X(1860)$  were also observed.
- ◆ Radiative decays  $\Upsilon(1S) \rightarrow \gamma X$  are analogous to that of  $J/\psi$ . The decay branching fraction is scaled down by  $1/25$ , due to the quark charges, masses and the total width of the quarkonia.
- ◆ CLEO II observed  $\Upsilon(1S) \rightarrow \gamma f_2(1270)$ , consistent with scaling down factor. CLEO also set an upper limit on  $f_J(2220)$  production.

# $\Upsilon(1S) \rightarrow \gamma h^+ h^-$



CLEO III

No significant structure at 2220 or 1860.

# $\Upsilon(1S) \rightarrow \gamma h^+ h^-$

- Confirm  $f_2(1270)$  in  $\pi^+\pi^-$  mode and establish J=2 assignment.

$$B(\Upsilon(1S) \rightarrow g f_2(1270)) = (10.2 \pm 0.8 \pm 0.7) \times 10^{-5}$$

- Observe  $f_2'(1525)$  in  $K^+K^-$  mode and establish J=2 assignment.

$$B(\Upsilon(1S) \rightarrow g f_2'(1525)) = (3.7_{-0.7}^{+0.9} \pm 0.8) \times 10^{-5}$$

- Set 90% CL upper limit on product branching fraction for  $f_J(2220)$ :


$$B(\Upsilon(1S) \rightarrow g f_J(2220)) \times B(f_J(2220) \rightarrow p^+ p^-) < 8 \times 10^{-7},$$

$$B(\Upsilon(1S) \rightarrow g f_J(2220)) \times B(f_J(2220) \rightarrow K^+ K^-) < 6 \times 10^{-7},$$

$$B(\Upsilon(1S) \rightarrow g f_J(2220)) \times B(f_J(2220) \rightarrow p \bar{p}) < 11 \times 10^{-7}.$$



- Set 90% CL upper limit on product branching fraction for X(1860):

$$B(\Upsilon(1S) \rightarrow g X(1860)) \times B(X(1860) \rightarrow p \bar{p}) < 5 \times 10^{-7}.$$



Published:  
PRD73, 032001(2006)

# $\Upsilon(1S) \rightarrow \gamma h^0 h^0$



10/31/06

- Measured  $f_2(1270)$  production rate in  $\pi^0\pi^0$  mode, consistent with  $\pi^+\pi^-$  mode.

$$B(\Upsilon(1S) \rightarrow g f_2(1270)) = (10.5 \pm 1.6_{-1.8}^{+1.9}) \times 10^{-5}$$

- No resonance structure for  $f_0(1500)$  and  $f_0(1710)$  and 90% upper limits are set. The limits are order of magnitude lower than QCD factorization calculation (PRD66, 074015):

$$B(\Upsilon(1S) \rightarrow g f_0(1500)) < 1.17 \times 10^{-5},$$

$$B(\Upsilon(1S) \rightarrow g f_0(1710)) \times B(f_0(1710) \rightarrow p^0 p^0) < 1.2 \times 10^{-6}.$$

- See two candidates in  $\eta\eta$  mode and no candidate in  $\pi^0\eta$  mode. UL are:

$$B(\Upsilon(1S) \rightarrow g f_0(1500)) \times B(f_0(1500) \rightarrow hh) < 3.0 \times 10^{-6},$$

$$B(\Upsilon(1S) \rightarrow g f_0(1710)) \times B(f_0(1710) \rightarrow hh) < 1.9 \times 10^{-6},$$


$$B(\Upsilon(1S) \rightarrow g p^0 h) < 2.8 \times 10^{-6}.$$

Submitted to PRD  
hep-ex/0512003

Jianchun Wang

12

# Search for $\Upsilon(1S) \rightarrow \gamma\eta, \gamma\eta'$ (I)



◆ Production rates of  $J/\psi \rightarrow \gamma\eta, \gamma\eta', \gamma f_2(1270)$  have been measured and

$$B(J/\psi \rightarrow gh')/B(J/\psi \rightarrow gf_2) \approx 3.4$$

$$B(J/\psi \rightarrow gh)/B(J/\psi \rightarrow gf_2) \approx 0.7$$


◆  $B(\Upsilon(1S) \rightarrow \gamma f_2(1270)) = 10.0 \times 10^{-5}$  by CLEO II and III.

◆ CLEO II set  $B(\Upsilon(1S) \rightarrow \gamma\eta') < 1.6 \times 10^{-5}$  and  $(\Upsilon(1S) \rightarrow \gamma\eta) < 2.1 \times 10^{-5}$ , corresponding to the ratio limits of 0.16 and 0.21 respectively.

◆ Several models (VDM, NRQCD, mixing with  $\eta_b$ ) try to explain the lower rates and need to be tested.


◆ Decay modes:  $\eta \rightarrow \gamma\gamma, \pi^+\pi^-\pi^0, \pi^0\pi^0\pi^0$  are searched. No candidate seen.

# Search for $\Upsilon(1S) \rightarrow \gamma\eta, \gamma\eta'$ (II)



**CLEO III  
Preliminary**

No Signal



$$\mathcal{B}(\Upsilon(1S) \rightarrow \gamma\eta) < 9.3 \times 10^{-7},$$

$$\mathcal{B}(\Upsilon(1S) \rightarrow \gamma\eta') < 1.77 \times 10^{-6}$$


Strongly disfavors mixing with  $\eta_b$ .

Still consistent with VDM.

Barely consistent with NRQCD.

# Search for $\Upsilon \rightarrow \gamma \mathfrak{R}$ , $\mathfrak{R} \rightarrow (\geq 4 h^\pm)$


Plot borrowed from  
PRD 74, 012003(2006)



- ◆ Two body decay with a narrow resonance  $\mathfrak{R}$  results in monochromatic  $\gamma$  in the lab frame.
- ◆ A bump above the smooth background indicates a narrow resonance.
- ◆ A series of fit at each  $E_\gamma$  step to Chebyshev polynomial function for background and Gaussian function for signal.
- ◆ The production upper limit at each  $E_\gamma$  is calculated from fit and map to  $M_{\mathfrak{R}}$ .

# Search for $\Upsilon \rightarrow \gamma \mathcal{R}, \mathcal{R} \rightarrow (\geq 4 h^\pm)$

CLEO III Preliminary  
hep-ex/0607054



	$B(\Upsilon \rightarrow \gamma \mathcal{R}, \mathcal{R} \rightarrow (\geq 4 h^\pm))$ 90%CL Upper Limit	
	All $M_{\mathcal{R}}$	$1.45 < M_{\mathcal{R}} < 5 \text{ GeV}$
$\Upsilon(1S)$	$1.05 \times 10^{-3}$	$1.82 \times 10^{-4}$
$\Upsilon(2S)$	$1.65 \times 10^{-3}$	$1.69 \times 10^{-4}$
$\Upsilon(3S)$	$2.47 \times 10^{-3}$	$3.00 \times 10^{-4}$



B.R upper limits are all  $\sim 10^{-4}$ .

No conflict with existing measurements




# Hadron Production in Gluon vs Quark

- ◆ Hadron productions in gluon rich and quark environment have been compared by many experiments.
- ◆ At LEP 3 jet ( $q\bar{q}g$ ) and 2 jet ( $q\bar{q}$ ) events were used for comparison.
- ◆ At CESR  $\Upsilon(1S) \rightarrow ggg$  and  $e^+e^- \rightarrow q\bar{q}$  were used. CLEO found that  $\phi$ ,  $\Lambda$  and  $p$  production rates are higher in  $ggg$  decays.
- ◆ Samples  $\Upsilon(1S) \rightarrow gg\gamma$  and  $e^+e^- \rightarrow q\bar{q}\gamma$  are better choices: Parton numbers are equal. Parton total energy can be equal.



# Comparison Between $gg\gamma$ and $q\bar{q}\gamma$




**CLEO III  
Preliminary**

Enhancement =

Number of hadron per  $gg\gamma$  event

Number of hadron per  $q\bar{q}\gamma$  event

# Comparison Between $ggg$ and $q\bar{q}$



Enhancement =

Number of hadron per  $ggg$  event

Number of hadron per  $q\bar{q}$  event

# Integrated Enhancement Comparison

1S		$gg\gamma$ vs. $q\bar{q}\gamma$	$ggg$ vs. $q\bar{q}$	1984 Study
Data	$\Lambda$	$1.86 \pm 0.25 \pm 0.03$	$2.668 \pm 0.027 \pm 0.051$	$\sim 3 \pm 0.3$
MC		$1.381 \pm 0.039$	$1.440 \pm 0.003$	
Data	$p$	$1.21 \pm 0.11 \pm 0.03$	$1.623 \pm 0.014 \pm 0.088$	$\sim 1.5 \pm 0.3$
MC		$1.582 \pm 0.034$	$1.331 \pm 0.005$	
Data	$\bar{p}$	$1.45 \pm 0.14 \pm 0.21$	$1.634 \pm 0.014 \pm 0.081$	
MC		$1.589 \pm 0.034$	$1.333 \pm 0.005$	
Data	$\phi$	$0.48 \pm 0.91 \pm 0.05$	$1.423 \pm 0.051 \pm 0.065$	$\sim 2 \pm 0.6$
MC		$0.702 \pm 0.027$	$0.836 \pm 0.003$	
Data	$f_2$	$1.34 \pm 0.84 \pm 0.05$	$0.658 \pm 0.029 \pm 0.065$	

CLEO III preliminary  
hep-ex/0607052

- ◆ CLEO III measurement of  $ggg$  vs  $q\bar{q}$  is consistent with CLEO II measurement.
- ◆ Enhancement effect is smaller in  $gg\gamma$  vs  $q\bar{q}\gamma$ .
- ◆ Enhancement is smaller for meson than baryon.

# Summary

- ◆ CLEO measured inclusive anti-deuteron and  $\eta'$  in  $\Upsilon$  decays.
- ◆ CLEO studied a group of radiative modes of  $\Upsilon$  decays.
- ◆ CLEO III dedicated  $\Upsilon(1S)$ ,  $\Upsilon(2S)$ ,  $\Upsilon(3S)$  data samples have produced many interesting results.
- ◆ They are rich laboratories to study hadronization process from gluon.