

Gauge Messenger Models

PRL 211803(2006) with R. Dermisek;
JHEP10(2006)001 with I.-W. Kim, and R. Dermisek;
hep-ph/0611xxx with K.-J. Bae, I. -W. Kim and R. Dermisek

Hyung Do Kim
Seoul National University

DPF JPS 2006
Hawaii, Oct. 31, 2006

Outline

- 1 Little Hierarchy Problem
- 2 Radiatively Generated Maximal Stop Mixing Scenario for Higgs Mass
- 3 Gauge Messenger Model
- 4 Neutralino Dark Matter in Gauge Messenger Model

- There are several good motivations for SUSY.
 - Stabilize weak scale (Higgs mass)
 - Gauge Coupling Unification with $M_{\text{GUT}} \sim 2 \times 10^{16}$ GeV.
 - Compatible with EW precision data.
 - Dark Matter with weak scale SUSY (with R-parity)

- SUSY suffers from *Little Hierarchy Problem*
 - We haven't found any experimental discovery of SUSY yet. Especially, LEP2 did not see **Higgs**. Higgs mass is logarithmically sensitive to the SUSY breaking scale.

$$m_h^2 \approx M_Z^2 \cos^2 2\beta + \frac{3G_F m_t^4}{\sqrt{2}\pi^2} \log \frac{m_{\tilde{t}}^2}{m_t^2}$$

- Current observational bound $m_h \geq 114\text{GeV}$. generically pushes up superpartner masses $m_{\tilde{t}}, m_{H_u}, \dots$ to $\mathcal{O}(\text{TeV})$
 M_Z is determined by soft SUSY breaking parameters.


$$\frac{M_Z^2}{2} \approx -\mu^2(M_Z) - m_{H_u}^2(M_Z)$$

- 0.5 % level fine-tuning → **Little Hierarchy Problem**

- Radiative correction to Higgs mass
 - Effective Potential has radiative corrections of the form

$$V^1(Q) = V^0(Q) + \Delta V^1(Q)$$

$$\Delta V^1(Q) = \frac{1}{64\pi^2} \text{Str} M^4(h) \left[\log \frac{M^2(h)}{Q^2} - \frac{3}{2} \right]$$



w/o mixing between \tilde{t}_L and \tilde{t}_R ,

$$m_h^2 \approx M_Z^2 \cos^2 2\beta + \frac{3G_F m_t^4}{\sqrt{2}\pi^2} \log \frac{m_{\tilde{t}}^2}{m_t^2}$$

→ **logarithmically** sensitive to $m_{\tilde{t}}^2$.

- To have $m_h > 114$ GeV, $m_{\tilde{t}} \sim \mathcal{O}(\text{TeV})$.

- Electroweak Symmetry Breaking is triggered mainly by two parameters μ and $m_{H_u}^2$ in MSSM:

$$\frac{M_Z^2}{2} \approx -\mu^2(M_Z) - m_{H_u}^2(M_Z)$$

- μ term does not change much from GUT scale value.

$$\frac{d\mu}{d \log Q} = \frac{\mu}{16\pi^2} (3y_t^2 + \dots) \propto \mu \quad (\text{no big change } 5 \sim 10\%)$$

- However, $m_{H_u}^2$ has a big radiative correction $\propto m_{\tilde{t}}^2$ through RG evolution.

$$\begin{aligned} \frac{dm_{H_u}^2}{d \log Q} &= \frac{3y_t^2}{8\pi^2} (m_{\tilde{Q}_3}^2 + m_{\tilde{t}^c}^2 + m_{H_u}^2 + |A_t|^2) \\ \delta m_{H_u}^2 &\approx -\frac{3y_t^2}{4\pi^2} m_{\tilde{t}}^2 \log \frac{M_{\text{GUT}}}{m_{\tilde{t}}} \approx -m_{\tilde{t}}^2 \sim \mathcal{O}(\text{TeV})^2 \end{aligned}$$

- We need fine-tuning of parameters to get $M_Z = 90$ GeV.

$$\frac{M_Z^2}{2} \approx -\mu^2(M_{\text{GUT}}) - m_{H_u}^2(M_{\text{GUT}}) + m_{\tilde{t}}^2$$

- For $m_{\tilde{t}} \sim 1$ TeV,
 $m_{H_u}^2(M_{\text{GUT}}) \sim (1 \text{ TeV})^2$
 and $\mu^2(M_{\text{GUT}}) \sim (1 \text{ TeV})^2$

$$\Delta = \frac{\frac{\delta M_Z^2}{M_Z^2}}{\frac{\delta M_{\text{UV}}^2}{M_{\text{UV}}^2}} \rightarrow \frac{2m_{H_u}^2(M_{\text{GUT}})}{M_Z^2} \sim 240$$

- 0.5% fine tuning.

Radiatively Generated Maximal Stop Mixing Scenario


with R.Dermisek, hep-ph/0601036 (PRL 211803(2006))

- Large Mixing between \tilde{t}_L and \tilde{t}_R helps higgs mass lift-up.


$$\mathcal{M}_{\tilde{t}}^2 = \begin{pmatrix} m_{\tilde{Q}_3}^2 + m_{\tilde{t}}^2 + \dots & -m_{\tilde{u}^c_3}(A_t^* + \mu \cot \beta) \\ -(A_t + \mu^* \cot \beta)m_{\tilde{u}^c_3}^* & m_{\tilde{u}^c_3}^2 + m_{\tilde{t}}^2 + \dots \end{pmatrix}$$

$$m_h^2 \sim M_Z^2 + \frac{3G_F m_t^4}{\sqrt{2}\pi^2} \left\{ \log \frac{M_S^2}{m_t^2} + \frac{A_t^2}{M_S^2} \left(1 - \frac{A_t^2}{12M_S^2} \right) \right\}$$

→ Maximum at $A_t = \pm\sqrt{6}M_S$.



- Physical Higgs Mass vs Stop Mixing (Carena et al)



- To satisfy LEP2 bound $m_h > 114\text{GeV}$ (using FeynHiggs)
for $m_{\tilde{t}}(M_Z) \approx 300\text{ GeV}$

$$|A_t(M_Z)| \approx 450\text{GeV}, \quad \tan \beta \gtrsim 50$$

$$|A_t(M_Z)| \approx 500\text{GeV}, \quad \tan \beta \gtrsim 8$$

$$|A_t(M_Z)| \approx 600\text{GeV}, \quad \tan \beta \gtrsim 6$$

- Therefore, $\left| \frac{A_t(M_Z)}{m_{\tilde{t}}(M_Z)} \right| \gtrsim 1.5$ is crucial.
- Unfortunately, the maximal mixing is not easily achieved due to the RG running.
- However, stop can be very light.


$$\log \frac{m_{\tilde{t}}^2}{m_t^2} = 3$$

$$\rightarrow m_{\tilde{t}_{(L,R)}} \simeq e^{-\frac{3}{2}} 1\text{ TeV} \simeq 250 \text{ or } 300\text{ GeV}.$$

$$\rightarrow m_{\tilde{t}_1} \sim m_{\tilde{t}} - m_t \sim 130\text{ GeV}. \quad \left(\frac{A_t}{m_{\tilde{t}}} = 2 \right)$$

- D0 squark gluino search (1st and 2nd generation)

- $m_{\tilde{g}} \geq 500$ GeV, $m_{\tilde{q}} \geq 200$ GeV
- $m_{\tilde{g}} \geq 230$ GeV, $m_{\tilde{q}} \geq 380$ GeV
- $m_{\tilde{g}} \geq 330$ GeV, $m_{\tilde{q}} \geq 330$ GeV



- Expressions for Weak scale parameters in terms of UV parameters
(for $\tan\beta = 10$)

$$m_{\tilde{t}}^2(M_Z) \approx 5.0M_3^2 + 0.6m_{\tilde{t}}^2 + 0.2A_t M_3$$

$$M_3(M_Z) \approx 3.0M_3$$

$$A_t(M_Z) \approx -2.3M_3 + 0.2A_t$$

$$\left| \frac{A_t(M_Z)}{m_{\tilde{t}}(M_Z)} \right| = \frac{|-2.3M_3 + 0.2A_t|}{\sqrt{5.0M_3^2 + 0.6m_{\tilde{t}}^2 + 0.2A_t M_3}} \lesssim 1 \text{ for positive } m_{\tilde{t}}^2.$$

- To achieve large stop mixing, we need **negative stop mass squared** at M_{GUT} .

- A_t at M_{GUT} does not help.

$$A_t(M_Z) \approx -2.3M_3 + 0.2A_t$$

$$\frac{dA_t}{d \log Q} = \frac{A_t}{16\pi^2} (18y_t^2 + \dots) \quad (\text{exponential suppression})$$

- Maximal mixing is not possible unless $A_t \geq 10M_3$ with opposite sign at the GUT scale.

- **Negative stop mass squared** also reduces fine-tuning.

- from the RG running of $m_{H_u}^2$,

$$\frac{dm_{H_u}^2}{d \log Q} \approx \frac{3y_t^2}{4\pi^2} m_t^2,$$

$m_{H_u}^2$ is lifted up by Yukawa loop if $m_{\tilde{t}}^2 < 0$. This enables $m_{H_u}^2$ to stay around M_Z^2 .

- In terms of fine-tuning to obtain M_Z^2 ,

$$M_Z^2 \approx -1.9\mu^2 - 1.2m_{H_u}^2 + 5.9M_3^2 + 1.5m_{\tilde{t}}^2 + \dots$$

→ For $m_{\tilde{t}}^2 \approx -4M_3^2$, stop contribution almost cancels gluino contribution so that μ and m_{H_u} can remain to be small.

- Near $m_{\tilde{t}}^2 \approx -4M_3^2$,

$$\left| \frac{A_t(M_Z)}{M_{\tilde{t}}(M_Z)} \right| = \frac{|-2.3M_3 + 0.2A_t|}{\sqrt{5.0M_3^2 + 0.6m_{\tilde{t}}^2 + 0.2A_tM_3}} \sim 1.5 - 0.2 \frac{A_t}{M_3}$$

- $A_t \sim -2.5M_3$ at the GUT scale gives $\frac{A_t}{m_{\tilde{t}}} = 2$ at the weak scale




FIG. 1: Renormalization group running of relevant SSBs for $\tan \beta = 10$ and GUT scale boundary conditions: $-A_t = M_3 = 200 \text{ GeV}$, $m_{\tilde{t}}^2 = -(400 \text{ GeV})^2$ and $m_{H_u}^2 = 0 \text{ GeV}^2$. In order to have both mass dimension one and two parameters on the same plot and keep information about signs, we define $m_{H_u} \equiv m_{H_u}^2 / \sqrt{|m_{H_u}^2|}$ and $m_{\tilde{t}} \equiv m_{\tilde{t}}^2 / \sqrt{|m_{\tilde{t}}^2|}$.

Cosmologically Viable?

- Large VEV Color/Charge Breaking Vacuum

- $m^2 < 0$ at high energy.
- Along the D-flat direction, there is no quartic coupling.
- Example


$$W = \frac{1}{M_{\text{Pl}}^3} (u^c d^c d^c)^2$$

$$V(\phi) = m^2 |\phi|^2 + \frac{1}{M_{\text{Pl}}^6} |\phi|^{10}.$$

- Then $\langle \phi \rangle \sim (m M_{\text{Pl}}^3)^{\frac{1}{4}} \sim 10^{14}$ GeV: Large VEV CCB minimum.
- The large VEV CCB minimum is always deeper than the EW vacuum (It is the AdS vacuum).

• Large VEV CCB Vacuum

- Effective potential is lifted up at finite temperature and there is no CCB vacuum. Universe will settle down to the EW vacuum after inflation. The tunneling rate to the deeper minimum is negligible.
- We can not reach AdS vacuum from expanding universe. It just collapses instantaneously.
- Coleman-De Luccia suppression



- EW Scale CCB Vacuum : Kusenko, Langacker, Segre
 - Large A_t can generate dangerous CCB vacuum at around the weak scale.
 - Assuming finite temperature after inflation, $T_R \geq 100$ GeV, the field is at the origin.
 - Phase transition to the EW vacuum is the second order while phase transition to A_t CCB vacuum is the first order.
 - The field just rolls to the EW vacuum unless tunneling rate to A_t CCB vacuum is very large.
 - Empirical bound :

$$|A_t|^2 + 3\mu^2 \leq 7.5(m_{\tilde{t}_L}^2 + m_{\tilde{t}_R}^2)$$

- Maximal stop mixing is compatible with the cosmological CCB bound.

(Middle) Conclusion

- Negative stop mass squared can make small μ and can make supersymmetry natural with maximal stop mixing. (other manifest solution : lower the cutoff Λ)
- Large A_t can be a solution of the little hierarchy problem.
- Either stop should be different from sleptons or gauginos should be non-universal (bino heavier than gluino).
- Naturalness points out the meta-stable vacuum.
- There are new regions of parameter space that should be seriously explored. (possibly better than CMSSM)
- But how?

Gauge Messenger Model

- In SUSY GUT, X, Y gauge bosons $\in SU(5)/G_{321}$ become massive at M_{GUT} by adjoint chiral superfield Σ . We consider the case where F -term of Σ is also induced.

$$SU(5) \xrightarrow{\langle \Sigma \rangle} G_{321} = SU(3) \times SU(2) \times U(1)$$

$$\Sigma = M_{\text{GUT}} \text{diag}(2, 2, 2, -3, -3) + \theta^2 F \text{diag}(2, 2, 2, -3, -3)$$

X, Y and $\lambda_{X,Y}$ are split in mass.

- $M_{\text{SUSY}} = \frac{\alpha_{\text{GUT}}}{4\pi} \left| \frac{F}{M} \right|$, b_G is β -func coeff. in $SU(5)$.
- $b_G = 3$ in the minimal case with a single adjoint.

Gauge Messenger Model

- Soft supersymmetry breaking parameters at M_{GUT} :

Gaugino

$$M_3 = -4M_{\text{SUSY}}, \quad M_2 = -6M_{\text{SUSY}}, \quad M_1 = -10M_{\text{SUSY}}$$

Scalar

$$m_{\tilde{Q}}^2 = (-20 + 3b_G)M_{\text{SUSY}}^2, \quad m_{\tilde{u}^c}^2 = (-16 + 4b_G)M_{\text{SUSY}}^2,$$

$$m_{\tilde{d}^c}^2 = (-12 + 2b_G)M_{\text{SUSY}}^2, \quad m_{\tilde{L}}^2 = (-12 + 3b_G)M_{\text{SUSY}}^2$$

$$m_{\tilde{e}^c}^2 = (-12 + 2b_G)M_{\text{SUSY}}^2, \quad m_{H_u}^2 = m_{H_d}^2 = (-12 + 3b_G)M_{\text{SUSY}}^2$$

Tri-linear

$$A_t = 10M_{\text{SUSY}}$$

- Gaugino Masses are **not universal** and have **opposite** sign to conventional GMSB.

$$\longrightarrow b_X \frac{\alpha}{4\pi} \frac{F}{M}$$

: Bino is the heaviest at M_{GUT} scale.


$$M_{\text{GUT}} \longrightarrow |M_1| : |M_2| : |M_3| = 2.5 : 1.5 : 1$$

$$M_Z \longrightarrow |M_1| : |M_2| : |M_3| \sim 1 : 1 : 2$$


- Negative soft scalar masses squared are generated and squark masses squared are most negative.
- $M_{\text{GUT}} : A_t$ is sizable ($A_t = -2.5M_3$).
- $M_Z : \frac{A_t(M_Z)}{m_{\tilde{t}}(M_Z)}$ close to maximal (~ 2).

- (Minimal) Gauge Messenger (using SoftSUSY)


- $M_{\text{SUSY}} = 56 \text{ GeV}$, $\tan \beta = 22$



- Gauge Messenger



- Pure Gauge Messenger : Parameter space



Geometric Hierarchy

- Dimopoulos, Raby

$$W = \text{Tr}\Sigma\tilde{\Sigma}^2 + \chi(\text{Tr}\tilde{\Sigma}^2 - M^2)$$

- F_Σ and F_χ can not be zero at the same time.
- M is the intermediate scale $\sim \sqrt{M_{\text{Pl}}M_Z}$.
- $F_\Sigma \sim F_\chi \sim M^2$.
- $\langle \Sigma \rangle \sim M_{\text{GUT}}$ becomes messenger scale.

Gravity Mediation

- Gravity mediation is not entirely negligible.

$$\frac{\alpha}{4\pi} \frac{F}{M_{\text{mess}}} \sim \frac{F}{M_{\text{Pl}}}$$

- Group theoretic numbers make gauge mediation dominates.

Additional effect to the soft parameters


- Adding a contribution from gravity mediation opens a possibility of a generating the μ term using the Giudice-Masiero mechanism. Comparable in size soft masses squared for H_u and H_d are also generated. We parametrize additional contribution to the Higgs soft masses squared by $c_{H_u} M_{\text{SUSY}}^2$ and $c_{H_d} M_{\text{SUSY}}^2$.
- We can add a universal contribution to all scalar masses from gravity mediation. We parametrize this contribution as $c_{\text{mSUGRA}} M_{\text{SUSY}}^2$.

Free parameters of gauge messenger model


- In the minimal ($b_G = 3$ and $N_{\text{mess}} = 0$) case, all the soft parameters at the GUT scale are determined by only one parameter M_{SUSY} . And at the EW scale, we need additional two parameters, μ and $\tan \beta$.
- In addition, we can include gravity contribution to Higgs sector : c_{H_u} and c_{H_d} .
- We can include universal gravity contribution to scalar masses : c_{mSUGRA} .
- Including these, every mass spectrum is determined by 6 parameters M_{SUSY} , μ , $\tan \beta$, c_{H_u} , c_{H_d} and c_{mSUGRA} .

- Gauge Messenger + Higgs



- $M_{\text{SUSY}} = 51 \text{ GeV}$, $\tan \beta = 32$, $c_{H_u} = 40$, $c_{H_d} = 34$,
 $m_{H_u}^2 = (-3 + c_{H_u})M_{\text{SUSY}}^2$



- Gauge Messenger + Higgs : Parameter space




Mass spectra for various parameter regions



- If $c_{m\text{SUGRA}} = 0$, NLSP is sneutrino (for low c_{H_u}) or stop (for high c_{H_u}) for low $\tan \beta$. And as $\tan \beta$ increases, down-type Yukawa couplings become large so that stau mixing becomes large and stau replaces sneutrino.
- But as $c_{m\text{SUGRA}}$ increases ($\gtrsim 5$), scalar masses become larger than gaugino masses, and the LSP is neutralino.

Mass spectra for various parameter regions



- In this figure, we can see that (N)LSP changes as $\tan \beta$ and c_{H_u} vary when $c_{\text{mSUGRA}} = 5$.

- Gravitino is LSP and $m_{\frac{3}{2}} \sim 100$ GeV
- Stau, sneutrino or stop is NSLP (Stau is a mixture of LR)
- In most of the parameter space, neutralino is LSP when $c_{\text{mSUGRA}} \geq 5$.

Neutralino masses in MSSM Lagrangian

where

$$N_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ 0 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

and

$$M_N^1 = \begin{pmatrix} M_1 & 0 & -M_Z s_{\beta+\frac{\pi}{4}} s_{\theta_W} & M_Z s_{\beta-\frac{\pi}{4}} s_{\theta_W} \\ 0 & M_2 & M_Z s_{\beta+\frac{\pi}{4}} c_{\theta_W} & -M_Z s_{\beta-\frac{\pi}{4}} c_{\theta_W} \\ -M_Z s_{\beta+\frac{\pi}{4}} s_{\theta_W} & M_Z s_{\beta+\frac{\pi}{4}} c_{\theta_W} & \mu & 0 \\ M_Z s_{\beta-\frac{\pi}{4}} s_{\theta_W} & -M_Z s_{\beta-\frac{\pi}{4}} c_{\theta_W} & 0 & -\mu \end{pmatrix}$$

Neutralino masses in MSSM Lagrangian

Since M_1 , M_2 and μ are sufficiently larger than M_Z , we can take off-diagonal terms as perturbation like quantum mechanics problems.

- When the difference between M_1 and M_2 is not extremely small, $(M_1 - M_2)\mu^2 \geq M_1 M_Z^2$, non degenerate perturbation formalism can be applied. First order result is given by,

$$N_{11} = 1, N_{12} = 0, N_{13} = \frac{-M_Z s_{\beta + \frac{\pi}{4}} s_{\theta_W}}{M_1 - \mu}, N_{14} = \frac{M_Z s_{\beta - \frac{\pi}{4}} s_{\theta_W}}{M_1 + \mu}, \quad (1)$$

where N_{1j} s are defined by lightest neutralino vector,

$$\chi_1 = N_{11} \tilde{B} + N_{12} \tilde{W}^0 + N_{13} \tilde{H}_1^0 + N_{14} \tilde{H}_2^0. \quad (2)$$

Neutralino masses in MSSM lagrangian

- As $N_{12} = 0$ at the first order, it is necessary to calculate the next order. Next order correction is small in general but can significantly alter the result when $M_1 \sim M_2$. The second order perturbation gives

$$\begin{aligned}
 N_{12} &\simeq \frac{M_Z^2 s_{\beta+\frac{\pi}{4}}^2 s_{2\theta_W}}{4\Delta M(\mu - M_1)} - \frac{M_Z^2 s_{\beta-\frac{\pi}{4}}^2 s_{2\theta_W}}{4\Delta M(\mu + M_1)} \\
 &\simeq \frac{M_Z^2 M_1 s_{2\theta_W}}{2\Delta M(\mu^2 - M_1^2)} = \frac{M_Z^2 s_{2\theta_W}}{2\epsilon(\mu^2 - M_1^2)},
 \end{aligned}$$

where $\Delta M = M_2 - M_1$ and $\epsilon = (M_2 - M_1)/M_1$. The second line is for $\tan \beta \geq 10$.

Neutralino in mSUGRA

- Since gaugino masses are universal at GUT scale in mSUGRA model, we know

$$|M_1| : |M_2| : |M_3| \sim 1 : 2 : 7$$

at EW scale from the relation $M_1/g_1^2 = M_2/g_2^2 = M_3/g_3^2$ at any scale.

- In mSUGRA, gaugino masses are a few hundred GeV, so $M_Z \ll |\mu \pm M_1|$. And since $\epsilon = 1$, the second order correction is negligible. Thus the lightest neutralino is mostly Bino.
- Mostly Bino neutralino gives too large relic density. It works only in **supercritically sensitive region** (co-annihilation or resonance cross section).
Arkani-Hamed, Delgado, Giudice (2006)

Main features in neutralino sector

- Starting from $|M_1|:|M_2|:|M_3|=2.5:1.5:1$ at the GUT scale, we get $|M_1|:|M_2|:|M_3|\sim 1:1:2$ at EW scale.
- Remind the second order term in χ eq.(15),

$$N_{12} \simeq \frac{M_Z^2 s_{2\theta_W}}{2\epsilon(\mu^2 - M_1^2)}.$$

Since $\epsilon \simeq 0.1$ in our case, the second order contribution becomes important and neutralino has sizable Wino contribution.

- Such a Wino-mixed neutralino can reduce the relic density and can explain the correct relic density with weak scale soft parameters.

Neutralino relic density for Bino/Wino, Higgsino mixture

- In most parameter region of mSUGRA, neutralino is usually pure Bino. And its relic density is much larger than observed value. If neutralino is a mixture of Bino and Wino, since the cross section of Wino is much larger than other contributions, from the eq.(44) relic density is approximately

$$\Omega_\chi h^2 \approx \frac{1}{|N_{12}|^2} \times 0.13 \left(\frac{M_2}{2.5 \text{TeV}} \right).$$

- If $N_{12} = 0.2$, i.e. 4% Wino, we get the Wino mass to fit the observed value,

$$0.09 \leq 25 \times 0.13 \left(\frac{M_2}{2.5 \text{TeV}} \right) \leq 0.13,$$

$$416 \text{GeV} \leq M_2 \leq 500 \text{GeV}$$

Bench mark point for relic density

- We choose the bench mark point as $M_{\text{SUSY}} = 90\text{GeV}$, $c_{\text{mSUGRA}} = 20$, $c_{H_u} = 10$, $c_{H_d} = -40$ and $\tan\beta = 10$.
- At EW scale, $M_1 = 379\text{GeV}$, $M_2 = 423\text{GeV}$, $\mu = -604\text{GeV}$, and

$$|N_{11}|^2 = 0.948, \quad |N_{12}|^2 = 0.034, \quad |N_{13}|^2 = 0, \quad |N_{14}|^2 = 0.018$$

$$\chi = N_{11}\tilde{B} + N_{12}\tilde{W}^0 + N_{13}\tilde{H}_1^0 + N_{14}\tilde{H}_2^0.$$



- Using the formulae for pure Wino and pure Higgsino cases, we can get the relic density of neutralino,

$$\begin{aligned} \Omega_\chi h^2 &\simeq \left(\frac{|N_{12}|^2}{\Omega_{\tilde{W}} h^2} + \frac{|N_{13}|^2 + |N_{14}|^2}{\Omega_{\tilde{H}} h^2} \right)^{-1} \\ &= \left[\frac{|N_{12}|^2}{0.13} \left(\frac{2.5\text{TeV}}{M_2} \right)^2 + \frac{|N_{13}|^2 + |N_{14}|^2}{0.10} \left(\frac{1.0\text{TeV}}{\mu} \right)^2 \right]^{-1} \\ &= 0.108 \end{aligned}$$

Bench mark point for relic density

- It satisfies the observed value, $0.9 \leq \Omega_{\chi} h^2 \leq 0.13$.
- In this parameter region, the numerical result is 0.088 for DARKSUSY.

Numerical results for the relic density



- The left-side figure is for simple gauge messenger case, i.e. no gravity contribution to Higgs. For large $\tan \beta$ region, co-annihilation with stau is shown and for small $\tan \beta$ region, pseudoscalar Higgs resonance appears.
- The right-side figure is for large gravity contribution to scalar and Higgs. There is bulk region to fit the observed value. There is no co-annihilation and no resonance.

Direct detection of neutralino

- Nucleon-neutralino scattering is described by effective four-fermi Lagrangian which is given by

$$\begin{aligned} \mathcal{L} = & \bar{\chi} \gamma^\mu \gamma^5 \chi \bar{q}_i \gamma^\mu (\alpha_{1i} + \alpha_{2i} \gamma^5) q_i + \alpha_{3i} \bar{\chi} \chi \bar{q}_i q_i \\ & + \alpha_{4i} \bar{\chi} \gamma^5 \chi \bar{q}_i \gamma^5 q_i + \alpha_{5i} \bar{\chi} \chi \bar{q}_i \gamma^5 q_i + \alpha_{6i} \bar{\chi} \gamma^5 \chi \bar{q}_i q_i, \end{aligned}$$



where $i = 1$ is for up-quark and $i = 2$ is for down-quark.

- For spin-independent case, α_{3i} term contributes. Since t-channel Higgs exchange scattering is dominant, cross section is approximately given by,

$$\sigma_{\chi p, n}^{\text{SI}} = \frac{g'^4 m_N^4}{4\pi} \left(\frac{\mu^2}{(\mu^2 - M_1^2)^2} \right) \left[\frac{X_d \tan \beta}{m_H^2} \left(\frac{M_1}{\mu} \right) + \frac{X_u}{m_h^2} \right]^2.$$

where X_u and X_d is determined by form factor of target materials (with decoupling approximation).

Direct detection of neutralino



Blue : spin-dependent, Red : spin-independent

- Our numerical result of nucleon-neutralino spin-independent cross section is below the current CDMS bound. It can be detected by CDMSII near future.

Conclusion

- **Negative Stop Mass Squared \rightarrow Meta-stable Vacuum**
 \rightarrow Large $A_t/M_{\tilde{t}}$ \rightarrow lightest Higgs mass bound even with light stop.
- **Non-universal Gaugino Masses**
 Gauge messenger model \rightarrow Non-universal gaugino mass, negative squark mass at GUT scale and large A term \rightarrow smaller fine-tuning
- **Degenerate Spectrum at the EW scale**
 Gluino and Bino/Wino (< 500 GeV) and squarks and sleptons also have similar masses (< 500 GeV).
- **Fine Tuning is improved up to 10 %**
 Almost factor 5 or 10 improvement compared to CMSSM.

Conclusion

- **Gravitino LSP and Stop NLSP**

Detection of NSLP decay using stopper might be possible.

- **Light Gluino, Light Stop**

Collider signal would be very interesting at the early year of LHC. (even at the Tevatron)

R-hadron (stop-light quark) is produced and is stable at the collider time scale

- **Neutralino Dark Matter**

Bino/Wino/Higgsino mixture with almost fixed ratio of mixing explains the dark matter relic density.

Direct detection might take longer than a few years.

- **mSUGRA vs Gauge Messenger Models**

Gauge messenger model is as good as mSUGRA or is better from phenomenological considerations.